PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A01N 51/00, 25/04, 25/30, 25/10

A1

(11) International Publication Number:

WO 98/34484

(43) International Publication Date:

13 August 1998 (13.08.98)

(21) International Application Number:

PCT/JP98/00522

(22) International Filing Date:

9 February 1998 (09.02.98)

(30) Priority Data:

9/26953

10 February 1997 (10.02.97) JP

(71) Applicant (for all designated States except US): TAKEDA CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka 541-0045 (JP).

(72) Inventors; and

- (75) Inventors'Applicants (for US only): KOIKE, Masahiko [JP/JP]; 7-9-1403, Kasuga 1-chome, Tsukuba-shi, Ibaraki 305-0821 (JP). SAKAI, Tomoyo [JP/JP]; 11-16-707, Sengen 2-chome, Tsukuba-shi, Ibaraki 305-0047 (JP).
- (74) Agents: ASAHINA, Tadao et al.; Osaka Plant of Takeda Chemical Industries, Ltd., 17-85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532-0024 (JP).

(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: AQUEOUS SUSPENSION OF AGROCHEMICAL

(57) Abstract

The present invention relates to an aqueous suspension comprising (i) a compound of formula (I) or a salt thereof, (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer. The aqueous suspension of the invention can be used with ad-

$$C1 \xrightarrow{H} CH_2 \xrightarrow{I} C = N - NO_2$$

$$CH_3 - N \xrightarrow{H} C = N - NO_2$$

$$CH_3 - N \xrightarrow{H} C = N - NO_2$$

$$CH_3 - N \xrightarrow{H} C = N - NO_2$$

$$CH_3 - N \xrightarrow{H} C = N - NO_2$$

$$CH_3 - N \xrightarrow{H} C = N - NO_2$$

$$CH_3 - N \xrightarrow{H} C = N - NO_2$$

vantage as a stable aqueous suspension of low viscosity providing for excellent delivery from a container, with excellent dispersibility in diluent water and excellent long-term fluidity free from caking due to precipitation of the suspended particles.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxem bourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad .
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	' Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU.	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DESCRIPTION

Aqueous Suspension of Agrochemical

5 [Technical Field]

The present invention relates to an agricultural aqueous suspension characterized by excellent pesticidal activity, low viscosity, and excellent suspension stability.

10

15

20

25

30

35

[Background Art]

Heretofore, as dosage forms for agricultural pesticides, dusts and granules have been primarily used. Those dosage forms are prepared by formulating a small amount of a pesticidally active substance with a large amount of a carrier such as clay, bentonite, calcium carbonate, etc. and, therefore, such formulations must be applied in quantities as large as 2 to 6 kilograms/10 ares in order to insure a uniform distribution of the active substance. This imposes a great burden on the farmers and, moreover, there are many disadvantages in transportation and storage. For this reason, so-called concentrates, which are supplied in small packages to provide for a greater ease of handling, are being developed.

Those concentrates are rich in the active substances and can be applied after dispersion in a suitable diluent, such as water, in the field. Examples include emulsifiable concentrates and wettable powders. However, emulsifiable concentrates require large quantities of an organic solvent for dilution, while it is unavoidable with wettable powders that preparation of sprays in the field involves contaminations with flying solid particles. For those reasons, the recent trend is toward refraining from using those formulations. Wettable granules, which are

20

30

35

granular versions of wettable powders have almost overcome the drawbacks of wettable powders but because the size of particles dispersed in water is as large as about 10 and odd μm , application of a dispersion of high concentration causes clogging of the spray nozzle, among other disadvantages.

The compound of the formula (I):

C1
$$\stackrel{\text{H}}{\downarrow}$$
 $\stackrel{\text{H}}{\downarrow}$ $\stackrel{\text{CH}_2-N}{\downarrow}$ $\stackrel{\text{C}}{\downarrow}$ $\stackrel{\text{C}}{$

or a salt thereof, is an agricultural pesticide having potent pest-controlling activity.

The demand exists today for development of a stable aqueous suspension of the compound (I) or a salt thereof which should have a low viscosity enough to permit smooth delivery from a container as well as satisfactory dispersibility in diluent water and longterm fluidity free from caking due to precipitation of the suspended particles.

25 [Disclosure of Invention]

The inventors of the present invention have made much research to overcome the above-mentioned disadvantages of the prior art and found that an aqueous suspension comprising (i) the compound (I) or a salt thereof, (ii) a condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer in combination may have a low viscosity of not more than 700 mPa·s with excellent dispersibility in diluent water and, in addition, have a long-term stability. The inventors

15

20

25

35

have made further research and have completed the present invention.

Namely, the present invention relates to:

(1) an aqueous suspension which comprises (i) a compound of the formula (I):

C1
$$N$$
 H CH_2-N $C=N-NO_2$ (1)

or a salt thereof, (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer,

- (2) the aqueous suspension as described in (1) above, wherein the aqueous suspension contains about 1 to 10 weight % of the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate in the total aqueous suspension,
- (3) the aqueous suspension as described in (1) above, wherein the absorptive water-soluble polymer is one or more (preferably one to three) polymers selected from the group consisting of polyvinylpyrrolidone, poly(alkylated vinylpyrrolidone), vinylpyrrolidonestyrene block copolymer and methyl vinyl ether-maleic anhydride copolymer,
- 30 (4) the aqueous suspension as described in (1) above, which further comprises an unabsorptive water-soluble polymer,
 - (5) the aqueous suspension as described in (4) above, wherein the unabsorptive water-soluble polymer is one or more (preferably one to three) polymers selected from the group consisting of a polymer in the polyol

series, a polysaccharides and polyethylene glycol, (6) the aqueous suspension as described in (4) above, wherein the unabsorptive water-soluble polymer is one or more (preferably one to three) polymers selected from the group consisting of xanthan gum, starch and polyethylene glycol,

- (7) the aqueous suspension as described in (1) above, which further comprises an auxiliary suspending agent,
- (8) the aqueous suspension as described in (7) above, wherein the auxiliary suspending agent is one or more (preferably one to three) substances selected from the group consisting of montmorillonite, colloidal silicon oxide, silicon oxide-aluminum oxide colloidal mixture and sepiolite,
- (9) a method for producing the aqueous suspension as described in (1) above, which comprises suspending (i) the compound of the formula (I):

20
$$C1 \qquad H \qquad H \qquad CH_2-N \qquad C=N-NO_2 \qquad (I)$$

$$CH_3-N \qquad C=N-NO_2 \qquad (I)$$

or a salt thereof, (ii) the condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, and (iii) the absorptive water-soluble polymer in water, and (10) a method for controlling a pest which comprises scattering the aqueous suspension as described in (1) above into a field.

[Best Mode for Carrying Out the Invention]

The compound (I) which is guanidine derivative for use in the aqueous suspension of the present invention occurs as <u>trans</u> (E) and <u>cis</u> (Z) isomers in respect of

the geometrical orientation of NO_2 . Preferred, however, is (E)-1-(2-chlorothiazol-5-ylmethyl)-3-methyl-2-nitroguanidine (hereinafter referred to briefly as compound (Ia)).

The salt of compound (I) includes agrochemically acceptable salts with inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, phosphoric acid, sulfuric acid, perchloric acid, etc. and salts with organic acids such as formic acid, acetic acid, tartaric acid, malic acid, citric acid, oxalic acid, succinic acid, benzoic acid, picric acid, p-toluenesulfonic acid, etc.

The production method for the compound (I) or a salt thereof is described in, <u>inter alia</u>, JP-A 157308/1991.

The proportion of compound (I) or a salt thereof in the aqueous suspension of the present invention is generally about 5-60 weight %, preferably about 5-50 weight %, and more preferably about 10-50 weight %, further more preferably 15-45 weight %.

The condensate of formaldehyde with aromatic sulfonic acid or a salt thereof which can be used in preparing the aqueous suspension of the present invention includes a condensate of formaldehyde with a sulfonic acid such as a phenylsulfonic acid, an arylsubstituted phenylsulfonic acid, a phenolsulfonic acid, an aryl-substituted phenolsulfonic acid, p-toluenesulfonic acid, a naphthalene sulfonic acid, an alkyl-substituted naphthalene sulfonic acid, an aralkyl-substituted naphthalene sulfonic acid, an anthracene sulfonic acid, or an alkyl-substituted anthracene sulfonic acid.

The aryl group of the aryl-substituted phenylsulfonic acid or aryl-substituted phenolsulfonic acid as mentioned above includes C_{6-14} aryl groups such as phenyl and naphthyl. The number of the substituents

5

10

15

20

25

30

35

10

15

is 1 to 3.

The alkyl group of the alkyl-substituted naphthalene sulfonic acid as mentioned above includes C_{1-6} alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl and tert-butyl. The number of the substituents is 1 to 7.

The aralkyl group of the aralkyl-substituted naphthalene sulfonic acid as mentioned above includes phenyl- C_{1-4} alkyl groups such as benzyl, phenethyl and phenylpropyl; benzhydryl; and trityl. The number of the substituents is 1 to 3.

The alkyl group of the alkyl-substituted anthracene sulfonic acid as mentioned above includes a C_{1-6} alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl and tert-butyl. The number of the substituents is 1 to 9.

The condensed degree between an aromatic sulfonic acid and formaldehyde is 1.2 to 30, preferably 1.2 to 10.

Among them as mentioned above, a naphthalene sulfonate-formaldehyde condensate is preferable.

The salt of the aromatic sulfonic acid includes alkaline metals such as sodium.

Specific examples of the condensate of
formaldehyde with aromatic sulfonic acid or a salt
thereof include New Kalgen FS-4 (sodium
naphthalenesulfonate-formaldehyde condensate; Takemoto
oil & fat co., Ltd.), New Kalgen 9131
(phenylphenolsulfonic acid-formaldehyde condensate;
Takemoto oil & fat co., Ltd.), Lavelin FM-L (sodium βnaphthalenesulfonate-formaldehyde condensate; Daiichi
Kogyo Seiyaku), New Kalgen WG-2 (sodium βnaphthalenesulfonate-formaldehyde condensate; Takemoto
oil & fat co., Ltd.), Demol N (sodium βnaphthalenesulfonate-formaldehyde condensate; Kao

Corporation), and Demol C (sodium aromaticsulfonate-

10

15

20

25

30

35

formaldehyde condensate; Kao Corporation).

Among them, especially New Kalgen FS-4 is preferable.

The polyoxyalkylene allyl phenyl ether sulfate includes those having a molecular weight of 500-5000.

The polyoxyalkylene in the polyoxyalkylene allyl phenyl ether sulfate includes polyoxyethylene, polyoxypropylene or polyoxybutylene. Especially, New Kalgen FS-7 (Takemoto oil & fat co., Ltd.) which contains mainly polyoxyethylene allyl phenyl ether sulfate is preferable.

The proportion of the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate in the total aqueous suspension of the present invention is generally about 1-10 weight %, preferably about 1-7 weight %.

One or more (preferable one to three) other surfactants such as nonionic or anionic surfactants besides the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, may be used in combination for preparing the aqueous suspension of the present invention.

The nonionic surfactant include a nonionic surfactant in the ether series such as polyoxyalkylene allyl phenyl ether, polyoxyethylene allyl phenyl etherformaldehyde condensate, polyoxyethylene alkyl phenyl ethers, polyoxyethylene nonyl phenyl ether, polyoxyethylene distyrenated phenyl ether, polyoxyethylene alkyl allyl ethers, polyoxyethylene alkyl allyl ethers, polyoxyethylene alkyl ethers, and polyoxyethylene-polyoxypropylene block copolymer. The nonionic surfactant that can be used in preparing the aqueous suspension of the invention is preferably one having an HLB number in the

20

25

30

35

range of 10-20. The more preferred HLB range is 12-18.

Among such surfactants, which are commercially available, are New Kalgen FS-1 (Takemoto oil & fat co., Ltd., polyoxyalkylene allyl phenyl ether), New Kalgen FS-11 (Takemoto oil & fat co., Ltd., polyoxyethylene allyl phenyl ether-formaldehyde condensate), Nonipol 100 (Sanyo Chemical Industries, LTD., polyoxyethylene nonyl phenyl ether), Noigen EA-177 (Daiichi Kogyo Seiyaku, polyoxyethylene distyrenated phenyl ether), Noigen EA-142 (Daiichi Kogyo Seiyaku, polyoxyethylene alkyl allyl ether), and Noigen EA-157 (Daiichi Kogyo Seiyaku, polyoxyethylene alkyl ether).

The anionic surfactant includes metal ligninsulfonates, polyoxyethylene allyl phenyl ether phosphate amine salts, polyoxyethylene alkyl phenyl ether phosphate amine salts, and sodium dialkylsulfosuccinates. To mention commercial products, there can be mentioned New Kalgen WG-4 (Takemoto oil & fat co., Ltd., metal ligninsulfonate), New Kalgen FS-3 (Takemoto oil & fat co., Ltd., polyoxyethylene allyl phenyl ether phosphate amine salt), and New Kalgen FS-31 and FS-32 (Takemoto oil & fat co., Ltd., polyoxyethylene alkyl phenyl ether phosphate amine salt).

The proportion of such a surfactant which may be used in combination in the total aqueous suspension of the present invention is generally about 1-10 weight % and preferably about 1-5 weight %.

To broaden the spectrum of utility of the aqueous suspension of the present invention, the suspension may be supplemented with one or more (preferably one to three) other agrochemicals. As such supplemental agrochemicals, insecticides, bactericides, etc. can be mentioned, which if such are solid at atmospheric temperature. The following are examples of such agrochemicals.

[Insecticides]

(I meet ele lues

15

20

25

Pyridafenthion, dimethoate, PMP, CVMP, dimethyl-vinphos, acephate, salithion, DEP, NAC, MTMC, MIPC, PHC, MPMC, XMC, bendiocarb, pirimicarb, mesomile, oxamyl, thiodicarb, cypermethrin, caltap hydrochloride, thiocyclam, bensultap, diflubenzuron, teflubenzuron, chlorfluazron, buprofezin, hexythiazox, phenbutatin oxide, pyridaben, clofentezine, nitenpyram, etc. [Bactericides]

Thiuram, captan, TPN, phthalide, trichlorophos methyl, phosethyl, methyl thiophanate, benomyl, carbendazole, thiabendazole, diethofencarb, iprodione, vinclozolin, procymidone, fluorimide, oxycarboxin, mepronil, flutolanil, bencyclane, metalaxyl, oxadixyl, triadimefon, hexaconazole, trifolin, blasticidin S, kasugamycin, polyoxin, validamycin A, mildéomycin, PCNB, hydroxyisoxazole, dazomet, dimethirimol, diclomezine, triazine, ferimzone, probenazole, isoprothiolane, tricyclazole, pyroquilone, oxolinic acid, etc.

The agrochemicals are not limited to the abovementioned ones but other suitable agrochemicals which are solid at atmospheric temperature can be added.

The absorptive water-soluble polymer is protective colloids which can absorb other hydrophobic substances by a hydrophobic group in the molecular, and which includes a polyvinylpyrrolidone, a poly(alkylated vinylpyrrolidone), a vinylpyrrolidone-styrene block copolymer and a methylvinyl ether-maleic anhydride copolymer.

The polyvinylpyrrodidone includes those having a mean molecular weight of 5,000-2,000,000 such as Agrimer 15 (mean molecular weight: about 8000), Agrimer 30 (mean molecular weight: about 57500), Agrimer 60 (mean molecular weight: about 406000) and Agrimer 90 (mean molecular weight: about 1270000), any of which are manufactured by ISP Ltd.

The alkyl of poly(alkylated vinylpyrrolidone) includes C_{1-30} alkyl groups such as methyl, ethyl, npropyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, pentyl, isopentyl, tert-pentyl, neopentyl, 5 hexyl, isohexyl, 2,2-dimethylbutyl, heptyl, 2methylhexyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, icosyl, henicosyl, dococyl, tricosyl and 10 tetracosyl. Among them, C_{1-24} alkyl groups are Especially, C_4 , C_{16} and C_{20} alkyl groups are preferable. Moreover, degree of alkylation of the poly preferable. (alkylated vinylpyrrolidone) is preferably 5-90%, more preferably 5-15% and 40-60%. The mean molecular weight 15 of the poly(alkylated vinylpyrrolidone) is preferably about 1,000 to about 100,000. Examples of the poly(alkylated vinylpyrrolidone) include Agrimer AL-10 (number of carbon atom in the alkyl group: 4, degree of alkylation: about 10%, mean molecular weight: about 20 70000), Agrimer AL-10LC (number of carbon atom in the alkyl group: 4, degree of alkylation: about 10%, mean molecular weight: about 70000), Agrimer AL-25 (number of carbon atom in the alkyl group: 16, degree of alkylation: about 50%, mean molecular weight: about 25 9500), and Agrimer AL-904 (number of carbon atom in the alkyl group: 4, degree of alkylation: about 10%, mean molecular weight: about 16,000), any of which are manufactured by ISP Ltd.

The vinylpyrrolidone-styrene block copolymer includes those having a mean molecular weight of about 40000 such as Agrimer ST (ISP Ltd.).

The methylvinylether-maleic anhydride copolymer includes free acid from having a mean molecular weight of about 100,000 to 300,000 such as Agrimer VEMA H-240 (ISP Ltd.), anhydride form having a mean molecular weight of about 100,000 to about 300,000 such as

30

35

10

20

25

30

Agrimer VEMA AN216 (ISP Ltd.), or half esters form having a mean molecular weight of about 20,000 to about 60,000 such as Agrimer ES-22 (ISP Ltd.).

As other absorptive water-soluble polymer which may be used in combination for preparing the aqueous suspension, poly(sodium acrylate) and carboxymethylcellulose are also mentioned.

As the absorptive water-soluble polymer, poly(alkylated vinylpyrrolidone) is preferable.

The proportion of the absorptive water-soluble polymer in the total aqueous suspension of the present invention is generally about 0.05-5 weight %, preferably about 0.1-3 weight %, more preferably about 0.1-2 weight %.

For the purpose of stabilizing and adjusting the viscosity of the suspension, the aqueous suspension of the present invention may be further supplemented with an unabsorptive water-soluble polymer and/or an auxiliary suspending agent.

The unabsorptive water-soluble polymer includes polymers in the polyol series (e.g. Agrisol FL-104FA, Kao Corporation); polysaccharides such as gum arabic, gelatin, guar gum, xanthan gum (e.g. Rhodopol 23; Rhone-Poulenc), starch (e.g. Stabirose MKK 100;

Matsutani Chemical Industry co., LTD.), starch derivatives such as hydrolyzed starch (e.g. Pine-Dex; Matsutani Chemical Industry co., LTD.), soluble-starch (e.g. Stabirose S-10; Matsutani Chemical Industry co., LTD.), enzymatically denatured starch (e.g. Amicol 6L; Nippon Starch Refining co., LTD.) and pregelatinized

starch (e.g. Cornalpha; Nippon Starch Refining co., LTD.) and dextrins (e.g. dextrin, cyclodextrin); and polyethylene glycol (e.g. PEG 6000; Sanyo Chemical Industries, LTD.).

Among them, the polysaccharides such as xanthan gum and starch, and polyethylene glycol are preferable.

20

When the unabsorptive water-soluble polymer is used, it is used in a proportion of generally about 0.05-10 weight %, preferably about 0.05-3 weight % based on the total suspension.

The auxiliary suspending agent is water-insoluble additive and includes montmorillonite (e.g. Kunipia F; Kunimine Industries co., LTD., Laponite; Nisshin Chemical Industries, LTD.), colloidal silicon oxide (e.g. Aerosil series such as Aerosil 200; Aerosil NIPPON), silicon oxide-aluminum oxide colloidal mixture (e.g. Aerosil COK84: SiO₂/Al₂O₃=82 to 86 (%)/14 to 18 (%); Aerosil NIPPON) and sepiolite (e.g. Aidplus ML70DS-NV; Mizusawa Industrial Chemicals Ltd.).

Among them, montmorillonite is preferable.

When the auxiliary suspending agent is used, it is used in proportion of generally about 0.05-10 weight %, preferably about 0.1-3 weight % based on the total suspension.

Those additives such as unabsorptive water-soluble polymers and auxiliary suspending agent can be used each alone or in combination.

In addition, other additives for routine use in aqueous suspensions in general can also be incorporated in the aqueous suspension of the present invention. For example, a pH control agent, an antifreeze agent, 25 an antifoam agent, and/or an antiseptic agent can be added liberally but selectively according to the amount of the agrochemical used. The pH control agent may for example be citric acid monohydrate. The antifreeze 30 agent that can be used includes, but is not limited to, ethylene glycol, diethylene glycol, and glycerin. antifoam agent may for example be a silicone series defoaming agent (e.g. Anti-foam E-20; KAO Corporation). The antiseptic agent includes, but is not limited to, 35 n-butyl p-hydroxybenzoate, potassium sorbate, and sorbic acid.

The pH control agent, when used, is added in a proportion of generally about 0.01-5 weight % or preferably about 0.05-3 weight % based on the total suspension. The antifreeze agent, when used, is used in a proportion of generally about 1-20 weight % or preferably about 3-10 weight % based on the total suspension. The antifoam agent, when used, is added in a proportion of generally about 0.05-0.5 weight % or preferably about 0.05-0.3 weight % based on the total suspension. The antiseptic agent, when used, is added in an amount of generally about 0.01-3 weight % or preferably about 0.01-1.5 weight % of the total suspension.

5

10

15

20

25

30

35

Q83448441 I

The aqueous suspension of the present invention can be produced by suspending (i) the compound (I) or a salt thereof, (ii) a condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer in water in accordance with the per se known method or any other technique analogous thereto. For example, there can be mentioned a process (1) which comprises mixing all the components together thoroughly in a high-speed mixer for about 20-60 minutes and pulverizing and dispersing the mixture by means of a wet mill such as Dynomill or a microfluidizer and a process (2) which comprises grinding bulk powders of the active substance, e.g. compound (I) or a salt thereof, in a dry mill such as Jetmizer and blending and dispersing the finely divided powders with the other components in a high-speed mixer for about 30-90 minutes.

As it may be understood from the above description of the production method, the effective ingredient for the aqueous suspension of the present invention is dispersed in finely divided form in water. In the presence of other powder components, too, the effective

10

15

20

25

ingredient is dispersed as efficiently in water. The mean diameter of the dispersed particles is not greater than about 10 μm , preferably 0.1-5 μm .

The crop plant to which the aqueous suspension of the present invention can be applied includes rice, wheat, barley, sugar beet, maize (Indian corn), cotton, vegetables (e.g. cabbage, Chinese cabbage, radish, cucumber, eggplant, potato, etc.), fruit-bearing trees (e.g. mandarin orange, peach, pear, etc.), tea, and tobacco.

The pest which can be controlled with the aqueous suspension of the present invention includes insects of the order <u>Hemiptera</u> {e.g. cabbage stink bug [<u>Eurydema</u> rugosum Motschulsky], black rice stink bug [Scotinophara lurida Burmeister], bean bug [Riptortus clavatus Thunberg], pear lace-bug [Stephanitis nashi Esaki et Takeya], brown planthopper [Nilaparvata lugens Stal], white-back planthopper [Sogota furcifera Horvath], smaller brown planthopper [Delphacodes striatella Fallén], green rice leafhopper [Nephotettix bipuntatus cincticeps Uhler], tea green leafhopper [Chlorita onukii Matsuda], greenhouse whitefly [Trialeurodes vaporariorum], arrow-head scale [Prontaspis yanonensis Kuwana], mulberry mealy bug [Pseudococcus comstocki Kuwana], soybean aphid [Aphis glycines Matsumura], turnip aphid [Rhopalosiphum pseudobrassicae Davis], cabbage aphid [Brevicoryne brassicae Linné], greenhouse aphid [Myzus persicae

Sulzer], cotton aphid [Aphis gossypii Glover], apple

aphid [Aphis pomi DeGeer], etc.}, the order Lepidoptera

{e.g. tabacco cutworm [Prodenia litura Fabricius],

diamond-back moth [Plutella maculipennis Curtis],

cabbage butterfly [Pieris rapae Linné], Asiatic rice

borer [Chilo suppressalis Walker], beet worm [Plusia

nigrisigna Walker], oriental tobacco budworm

[Helicoverpa assulta assulta Guenée], armyworm

[<u>Leucania separata</u> Walker], cabbage armyworm [<u>Mamestra</u> brassicae Linné], smaller tea tortrix [Adoxophyes orana Fischer von Röslerstamm] (apple), cotton leaf roller [Sylepta derogata Fabricius], grass leaf roller 5 [Cnaphalocrocic medinalis Guénée], potato tuberworm [Phthorimaea operculella Zeller], tea leaf roller [Caloptilia theivora Walsingham], smaller tea tortrix [Adoxophyes orana Fischer von Röslerstamm] (tea), etc.}, the order Coleoptera {e.g. potato lady beetle 10 [Epilachna sparsa orientalis Dieke], cucurbit leaf beetle [Rhaphidopalpa femoralis Motschulsky], striped cabbage flea-beetle [Phyllotreta vittata Fabricius], Colorado potato bug [Leptinotarsa decemlineata], beet tortoise beetle [Cassida nebulosa Linné], rice leaf 15 beetle [Oulema oryzae Kuwayama], rice plant weevil [Echinocnemus squameus Billberg], vegetable weevil [Listroderes obliquus Klug], etc.}, the order Diptera {e.g. oriental house fly [Musca domestica vicina Macquart], pale house mosquito [Culex pipiens pallens 20 Coquillett], common horse fly [Tabanus trigonus Coquillett], onion maggot [Hylemya antiqua Meigen], seed-corn maggot [Hylemya platura Meigen], etc.}, the order Orthoptera {e.g. Asiatic locust [Locusta migratoria Linné], African mole cricket [Gryllotalpa 25 africana Palisot de Beauvois], etc.}, reticulopterous insects {e.g. German cockroach [Blattella germanica Linné], smoky-brown cockroach [Periplaneta fuliginosa Serville], etc.}, spider mites {e.g. two-spotted spider mite [Tetranychus urticae Koch], citrus red mite 30 [Panonychus citri McGregor], Kanzawa spider mite [Tetranychus kanzawai Kishida], carmine mite [Tetranychus telarius Linné], European red mite [Panonychus ulmi Koch], Japanese citrus rust mite [Aculus pelekassi Keifer], etc.}, nematodes {e.g. rice 35 white-tip nematode [Aphelenchoides besseyi Christie] etc.} and so on.

10

20

The aqueous suspension of the present invention is a safe and useful agrochemical composition with low toxicity against plants and mammals. This agrochemical composition can be used either as it is or as dilution in a suitable medium, e.g. water, when used. recommended method for application depends on the crop plant applied, but is generally scattering by means of spraying or dropping, or dipping to field, i.e. paddy field, plow-land, orchard, grass-land or non-crop land in accordance with the per se known method. specific method for application includes aerial broadcast, soil treatment, folial application, nursery box treatment, hand application along rows, seed dressing, bed soil treatment, directed application to 15 infesting pests, application to paddy flood water, etc.

The recommended dose of the aqueous suspension of the present invention depends on the concentration of the agrochemical active ingredient but is generally about 1-500 g, preferably about 1-200 g, as active ingredient, per 10 ares for paddy field and plow-land (growing tea, wheat, sugar beet, maize, potato, cotton, etc.) pests and orchard pests.

[Examples]

25 The following examples and test examples are intended to illustrate the present invention in further The compound (Ia) used here was prepared in accordance with the procedure described in JP-A 157308/1991 (Table 4; Compound No. 19). It should be 30 understood that all parts and percents (%) are by weight unless otherwise indicated.

[Example 1]

	Compound (Ia)	20	parts
	New Kalgen FS-4	5	parts
35	Agrimer 30	1	part
	Polyethylene glycol (PEG6000)	2	parts

Rhodopol 23 0.1 part Anti-foam E-20 0.2 part Ethylene glycol 4 parts Kunipia F 1 part 5 n-Butyl p-hydroxybenzoate 0.1 part Water add 100 parts The above components were thoroughly mixed in a high-speed mixer and wet-pulverized (one pass) by means of Dynomill (Shinmaru Enterprises, 1.0 mm glass beads, 10 packing rate 80%, peripheral speed 15 m/s) to provide an aqueous suspension containing 20% of compound (Ia). The term of "add 100 parts" means to make 100 parts of aqueous suspension by adding water besides other components. 15 [Example 2] Compound (Ia) 20 parts New Kalgen FS-4 2 parts Agrimer AL-10 1 part Stavirose MKK 100 0.5 part 20 Anti-foam E-20 0.2 part Ethylene glycol 4 parts Aerosil COK84 1 part n-Butyl p-hydroxybenzoate 0.1 part Water add 100 parts 25 Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 20% of compound (Ia). [Example 3] Compound (Ia) 20 parts 30 New Kalgen FS-7 3 parts Agrimer ST 1 part Anti-foam E-20 0.2 part Ethylene glycol 4 parts Aidplus ML70DS-NV 1 part 35 0.1 part n-Butyl p-hydroxybenzoate

add 100 parts

Water

Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 20% of compound (Ia).

[Example 4]

5	Compound (Ia)	20 parts
	New Kalgen FS-7	3 parts
	Agrimer VEMA H-240	0.5 part
	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
10	Kunipia F	1.5 part
	Citric acid monohydrate	0.1 part
	n-Butyl p-hydroxybenzoate	0.1 part
•	Water add	100 parts
	Using the above components	, the procedure of
15	Example 1 was repeated to provi	de an aqueous suspension
·	containing 20% of compound (Ia)	•

[Example 5]

•	Compound (Ia)		40	parts
	New Kalgen FS-7		5	parts
20	Agrimer AL-30		1	part
	Anti-foam E-20		0.	2 part
	Ethylene glycol		7	parts
	Aerosil COK84		1.	5 part
	Rhodopol 23		0.	2 part
25	n-Butyl p-Hydroxybenzo	ate	0.	1 part
	Water	add	100	parts

Using the above components, the procedure of Example 1 was repeated to provide an aqueous suspension containing 40% of compound (Ia).

30 [Example 6]

	Compound (Ia)	21 parts
	New Kalgen FS-4	2 parts
	Agrimer AL-10	0.5 part
	Stabirose MKK 100	0.3 part
35	Rhodopol 23	0.1 part
	Kunipia F	0.8 part

	n-Butyl p-hydroxybenzoate	0.1 part
٠	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
	Water add	100 parts
5	Using the above components,	the procedure of
	Example 1 was repeated to provide	e an aqueous suspension
	containing 21% of compound (Ia).	
	[Example 7]	
	Compound (Ia)	21 parts
10	New Kalgen FS-4	2 parts
	Agrimer AL-10	0.5 part
	Polyethylene glycol (PEG600)	0) 2.0 part
	Rhodopol 23	0.05 part
	Kunipia F	0.65 part
15	n-Butyl p-hydroxybenzoate	0.1 part
	Anti-foam E-20	0.2 part
	Ethylene glycol	4 parts
	Water add	d 100 parts
	Using the above components,	the procedure of
20	Example 1 was repeated to provide	e an aqueous suspension
	containing 21% of compound (Ia).	
	[Example 8]	
	Compound (Ia)	21 parts
	New Kalgen FS-4	2 parts
25	Agrimer AL-10	0.5 part
	Polyethylene glycol (PEG600	0) 2 parts
	Kunipia F	0.8 part
	n-Butyl p-hydroxybenzoate	0.1 part
	Anti-foam E-20	0.2 part
30	Ethylene glycol	4 parts
	Water ad	d 100 parts
	Using the above components,	the procedure of
	Example 1 was repeated to provid	e an aqueous suspension
	containing 21% of compound (Ia).	
35	[Test Example 1]	
	Using the aqueous suspension	ns prepared in Examples

10

15

20

25

1-8, the viscosity, water dispersibility, stability of 10-fold diluted suspension and tendency toward caking of each aqueous suspension were evaluated by the following methods.

(1) Viscosity

Viscosity was measured with a Type B viscosimeter at 25°C.

(2) Water dispersibility

A hollow cylinder of 500 ml capacity was filled in with 500 ml of water. Then, 0.5 ml of the test suspension was added dropwise and the dispersion was evaluated according to the criteria: (A) excellent, (B) good, (C) poor, and (D) not dispersed.

(3) Stability test of 10-fold diluted suspension

A hollow cylinder of 500 ml capacity was filled in with 450 ml of water and 50 ml of diluted suspension (flowable), and then turned upside down 10 times.

The test suspensions were respectively allowed to stand for 15 hours and the degree of the separation of water-phase was investigated.

(4) Tendency toward caking

The test suspensions were respectively allowed to stand at room temperature for 12 months following preparation and the degree of caking on the container bottom was investigated.

The results are shown in Table 1. [Table 1]

30	Sample	Viscosity (mPa·s)	Water disper- sibility	Degree of separation (%)	Caking
	Ex. 1	230	A	2.1	non
	Ex. 2	150	A	1.8	non
	Ex. 3	65	A	3.2	non
35	Ex. 4	370	A	3.0	non
	Ex. 5	280	A	1.1	non
فمسد	Ex. 6	290	A	2.1	non
	Ex. 7	390	A	2.4	non
	Ex. 8	250	А	2.1	non

It is clear from Table 1 that the aqueous suspension of the present invention is of low viscosity, i.e. not higher than 700 mPa·s, with good water dispersibility and has a good shelf-life (stability of the stock suspension).

[Industrial Applicability]

The aqueous suspension comprising (i) the compound (I) or a salt thereof, (ii) the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer can be used with advantage as a highly stable agrochemical preparation with a low viscosity, i.e. not higher than 700 mPa·s, thus providing for excellent delivery from a container with excellent dispersibility in diluent water and excellent long-term fluidity free from caking due to precipitation of the suspended particles.

CLAIMS

1. An aqueous suspension which comprises (i) a compound of the formula (I):

$$\begin{array}{c|c}
C1 & H \\
S & CH_2 - N \\
CH_3 - N \\
H
\end{array}$$

$$C = N - NO_2 \qquad (I)$$

or a salt thereof, (ii) a condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or a polyoxyalkylene allyl phenyl ether sulfate, and (iii) an absorptive water-soluble polymer.

- 2. The aqueous suspension as claimed in claim 1 wherein the aqueous suspension contains about 1 to 10 weight % of the condensate of formaldehyde with aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate in the total aqueous suspension.
- 3. The aqueous suspension as claimed in claim 1 wherein the absorptive water-soluble polymer is one or more polymers selected from the group consisting of polyvinylpyrrolidone, poly(alkylated vinylpyrrolidone), vinylpyrrolidone-styrene block copolymer and methyl vinyl ether-maleic anhydride copolymer.
- 4. The aqueous suspension as claimed in claim 1 which further comprises an unabsorptive water-soluble polymer.
- 5. The aqueous suspension as claimed in claim 4 wherein the unabsorptive water-soluble polymer is one or more polymers selected from the group consisting of a polymer in the polyol series, a polysaccharide and polyethylene glycol.
- 6. The aqueous suspension as claimed in claim 4 wherein

the unabsorptive water-soluble polymer is one or more polymer selected from the group consisting of xanthan gum, starch and polyethylene glycol.

23

- 7. The aqueous suspension as claimed in claim 1 which further comprises an auxiliary suspending agent.
- 8. The aqueous suspension as claimed in claim 7 wherein the auxiliary suspending agent is one or more substances selected from the group consisting of montmorillonite, colloidal silicon oxide, silicon oxide-aluminum oxide colloidal mixture and sepiolite.
- 9. A method for producing the aqueous suspension as claimed in claim 1 which comprises suspending (i) the compound of the formula (I):

$$C1 \longrightarrow H$$

$$CH_2 \longrightarrow N$$

$$CH_3 \longrightarrow H$$

$$C = N \longrightarrow NO_2$$

$$C = N \longrightarrow NO_2$$

$$C = N \longrightarrow NO_2$$

$$(1)$$

or a salt thereof, (ii) the condensate of formaldehyde with an aromatic sulfonic acid or a salt thereof or the polyoxyalkylene allyl phenyl ether sulfate, and (iii) the absorptive water-soluble polymer in water.

10. A method for controlling a pest which comprises scattering the aqueous suspension as claimed in claim 1 into a field.

INTERNATIONAL SEARCH REPORT

Int. ational Application No PCT/JP 98/00522

A. CLASSI	FICATION OF SUBJECT MATTER		
IPC 6	A01N51/00 A01N25/04 A01N25	6/30 A01N25/10	e e e e e e e e e e e e e e e e e e e
			!
According to	o International Patent Classification(IPC) or to both national class	sification and IPC	
	SEARCHED		
	ocumentation searched (classification system followed by classific	cation symbols)	
IPC 6	AOIN		
Documenta	tion searched other than minimum documentation to the extent the	at such documents are included in the fields sea	arched
		, and the second	
Clarencia d	the base seem that during the international accords (name of date	has and when practical ageral terms upod	
Flectionic	lata base consulted during the international search (name of data	t base and, where practical, search terms used)	
C DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the	relevant passages	Relevant to claim No.
Α	EP 0 418 199 A (CIBA GEIGY AG)	20 March	1-10
	1991		
	see page 2, line 1 - line 38	lino 21	
	see page 10, line 35 - page 12 see pages 14-30, table 1, compo		
	and pages 32, formulation examp		
A	EP 0 425 978 A (AGRO KANESHO CO	1-10	
	see page 3, line 1 - line 34		
	see page 7, table 1, compound 8		
	see page 7, line 54 - page 8,	line 6	
	see page 10, formulation examp	ies 1-3	
		-/	
ļ			
X Fun	ther documents are listed in the continuation of box C.	Patent family members are tisted	in annex.
° Special c	ategories of cited documents :	"T" later document published after the inte	ernational filing date
	ent defining the general state of the art which is not	or priority date and not in conflict with cited to understand the principle or the	the application but
"E" earlier	dered to be of particular relevance document but published on or after the international	invention "X" document of particular relevance; the	claimed invention
filing	date ent which may throw doubts on priority claim(s) or	cannot be considered novel or canno involve an inventive step when the de	t be considered to
which	n is cited to establish the publication date of another on or other special reason (as specified)	"Y" document of particular relevance; the cannot be considered to involve an ir	claimed invention
"O" docum	nent referring to an oral disclosure, use, exhibition or means	document is combined with one or ments, such combination being obvio	ore other such docu-
"P" docum	nent published prior to the international filing date but	in the art. "&" document member of the same paten	•
	than the priority date claimed actual completion of theinternational search	Date of mailing of the international se	
			·
1	15 June 1998	30/06/1998	
Name and	mailing address of the ISA	Authorized officer	
	European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk		
	Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Muellners, W	

Form PCT/ISA/210 (second sheet) (July 1992)

2

INTERNATIONAL SEARCH REPORT



	lation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category '	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL PATENTS INDEX, DOCUMENTATION ABSTRACTS JOURNAL Week 9022 25 July 1990 Derwent Publications Ltd., London, GB; AN 90169074 XP002067921 & JP 02 111 703 A (DAIICHI KOGYO SEIYAKU) , 24 April 1990 see abstract	1-10
Α	EP 0 425 729 A (NIHON NOHYAKU CO LTD) 8 May 1991 see page 2, column 23 - page 3, column 20 see pages 6-9, table 1, comparative example 9 see page 10, line 1 - line 31	1-10
Α	CENTRAL PATENTS INDEX, BASIC ABSTRACTS JOURNAL Week 8335 1986 Derwent Publications Ltd., London, GB; AN 83-750619 XP002067922 & JP 58 124 702 A (KUMIAI CHEM IND KK), 25 July 1983 see abstract	1-10
A .	CHEMICAL PATENTS INDEX, DOCUMENTATION ABSTRACTS JOURNAL Week 9301 10 March 1993 Derwent Publications Ltd., London, GB; AN 93-005486 XP002067923 & JP 04 334 306 A (NIHON NOYAKU CO LTD) , 20 November 1992 see abstract	1-10
. · ·		

2

INTERN ONAL SEARCH REPORT

information on patent family members

.tional Application No PCT/JP 98/00522

Patent documentcited_in_search_report		Publication date	Patent familymember(s)		Publication	
EP 0418199	Α	20-03-1991	AU	6245690 A	21-03-1991	
			BG	60863 B	31-05-1996	
			CA	2025072 A	14-03-1991	
			CN	1050186 A	27-03-1991	
			EG	19741 A	31-01-1996	
			JP .	3109374 A	09-05-1991	
			PT	95275 A	22-05-1991	
EP 0425978	Α	08-05-1991	JP	3200768 A	02-09-1991	
			AU	6396490 A	30-05-1991	
			CN	1051728 A	29-05-1991	
			US	5166164 A	24-11-1992	
EP 0425729	A	08-05-1991	JP	1283201 A	14-11-1989	
			KR	9506924 B	26-06-1995	